

Inequalities for generalized parton distributions H and E

P.V. Pobylitsa

Institute for Theoretical Physics II, Ruhr University Bochum,
D-44780 Bochum, Germany
Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg, 188350, Russia

Abstract

New positivity bounds are derived for quark and gluon generalized parton distributions H and E in spin 1/2 hadrons.

Generalized parton distributions (GPDs) appear in the QCD description of a number of hard exclusive processes [1, 2, 3, 4, 5]. Since our current knowledge about GPDs is rather poor, any additional information deserves attention and from this point of view so called positivity bounds [6, 7, 8] for GPDs are of certain interest.

In this paper the following bound for quark generalized parton distributions H_q and E_q of spin 1/2 hadrons is derived

$$\left| H_q(x, \xi, t) - \frac{\xi^2}{1 - \xi^2} E_q(x, \xi, t) \right|^2 + \left| \frac{\sqrt{t_0 - t}}{2M\sqrt{1 - \xi^2}} E_q(x, \xi, t) \right|^2 \leq \frac{q(x_1)q(x_2)}{1 - \xi^2}. \quad (1)$$

Here the notation of X. Ji is used for quark GPDs H_q, E_q and for their arguments x, ξ, t (see e.g. [9]), M is the mass of the hadron. The usual (forward) unpolarized quark distributions $q(x)$ are taken at values of x

$$x_1 = \frac{x + \xi}{1 + \xi}, \quad x_2 = \frac{x - \xi}{1 - \xi}. \quad (2)$$

Inequality (1) is written for the region $|\xi| < |x|$. The parameter

$$t_0 = -\frac{4\xi^2 M^2}{1 - \xi^2} \quad (3)$$

corresponds to the maximal value of the squared momentum transfer t ($t \leq t_0 \leq 0$).

Obviously the bound (1) is stronger than the following inequality derived in ref. [10]

$$\left| H_q(x, \xi, t) - \frac{\xi^2}{1 - \xi^2} E_q(x, \xi, t) \right| \leq \sqrt{\frac{q(x_1)q(x_2)}{1 - \xi^2}}, \quad (4)$$

We also see from inequality (1) that

$$\frac{\sqrt{t_0 - t}}{2M} |E_q(x, \xi, t)| \leq \sqrt{q(x_1)q(x_2)}. \quad (5)$$

This inequality agrees with the one derived in ref. [10] up to a typo there [11].

As noticed in ref. [10] the earlier bound suggested in refs. [7, 8]

$$|H_q(x, \xi, t)| \leq \sqrt{\frac{q(x_1)q(x_2)}{1 - \xi^2}} \quad (6)$$

is not justified since in its derivation the E_q contribution was overlooked.

The derivation of inequality (1) starts from the positivity condition [7, 8, 10]

$$\left\| \sum_{k=1}^2 b_\lambda^k \int \frac{d\tau}{2\pi} e^{i\tau x_k(P_k n)} \psi_\alpha(\tau n) |P_k, \lambda\rangle \right\|^2 \geq 0. \quad (7)$$

The lhs contains a superposition of two states obtained by acting with quark fields ψ on a nucleon state $|P_k, \lambda\rangle$ with momentum P_k and polarization λ ; n is a light-cone vector and the coefficients b_λ^k are arbitrary. Expanding this squared sum and summing over the “good” spin components α of the quark fields we obtain

$$b_\lambda^{1*} b_\lambda^1 q(x_1) + b_\lambda^{2*} b_\lambda^2 q(x_2) + \sqrt{1 - \xi^2} [b_\lambda^{1*} \mathcal{H}_{\lambda', \lambda}^q(-\Delta) b_\lambda^2 + b_\lambda^{2*} \mathcal{H}_{\lambda', \lambda}^q(\Delta) b_\lambda^1] \geq 0. \quad (8)$$

The summation over repeated indices is implied, $\Delta = P_2 - P_1$, and the matrix $\mathcal{H}_{\lambda', \lambda}^q$ is taken from ref. [10]

$$\mathcal{H}_{\lambda', \lambda}^q(\Delta) = \begin{pmatrix} H_q - \frac{\xi^2}{1 - \xi^2} E_q & \frac{-\Delta^1 + i\Delta^2}{2M(1 - \xi^2)} E_q \\ \frac{\Delta^1 + i\Delta^2}{2M(1 - \xi^2)} E_q & H_q - \frac{\xi^2}{1 - \xi^2} E_q \end{pmatrix}_{\lambda', \lambda}. \quad (9)$$

The inequality (8) should hold for any b_λ^k . Taking special cases $b_-^1 = b_-^2 = 0$ or $b_-^1 = b_+^2 = 0$ one reproduces the bounds (4), (5).

Now let us consider the case of arbitrary b_λ^k . One can rewrite inequality (8) in the following form

$$\begin{aligned} & \frac{1}{\sqrt{1 - \xi^2}} [b_\lambda^{1*} b_\lambda^1 q(x_1) + b_\lambda^{2*} b_\lambda^2 q(x_2)] + (b_\lambda^{1*} b_\lambda^2 + b_\lambda^{2*} b_\lambda^1) \left(H_q - \frac{\xi^2}{1 - \xi^2} E_q \right) \\ & + i (b_{\lambda_2}^{1*} b_{\lambda_1}^2 - b_{\lambda_2}^{2*} b_{\lambda_1}^1) (\Delta^1 \sigma^2 - \Delta^2 \sigma^1)_{\lambda_2 \lambda_1} \frac{E_q}{2M(1 - \xi^2)} \geq 0 \end{aligned} \quad (10)$$

where σ^k are Pauli matrices. Rotating the spinor indices λ of b_λ^k by an $SU(2)$ transformation one can diagonalize $\Delta^1 \sigma^2 - \Delta^2 \sigma^1 \rightarrow |\Delta_\perp| \sigma^3$ so that the inequality takes the form

$$\frac{1}{\sqrt{1 - \xi^2}} [b_\lambda^{1*} b_\lambda^1 q(x_1) + b_\lambda^{2*} b_\lambda^2 q(x_2)]$$

$$\begin{aligned}
& + (b_{\lambda}^{1*} b_{\lambda}^2 + b_{\lambda}^{2*} b_{\lambda}^1) \left(H_q - \frac{\xi^2}{1 - \xi^2} E_q \right) \\
& + i (b_{\lambda_2}^{1*} b_{\lambda_1}^2 - b_{\lambda_2}^{2*} b_{\lambda_1}^1) |\Delta_{\perp}| \sigma_{\lambda_2 \lambda_1}^3 \frac{E_q}{2M(1 - \xi^2)} \geq 0.
\end{aligned} \tag{11}$$

We see that we have two independent inequalities for b_+^k and b_-^k

$$\sum_{k,k'} b_{\pm}^{k'} \begin{pmatrix} \frac{q(x_1)}{\sqrt{1 - \xi^2}} & H_q - \frac{\xi^2}{1 - \xi^2} E_q \pm i \frac{|\Delta_{\perp}| E_q}{2M(1 - \xi^2)} \\ H_q - \frac{\xi^2}{1 - \xi^2} E_q \mp i \frac{|\Delta_{\perp}| E_q}{2M(1 - \xi^2)} & \frac{q(x_2)}{\sqrt{1 - \xi^2}} \end{pmatrix}_{k'k} b_{\pm}^k \geq 0. \tag{12}$$

The positivity of this quadratic form immediately leads us to the constraint

$$\left(H_q - \frac{\xi^2}{1 - \xi^2} E_q \right)^2 + \left(|\Delta_{\perp}| \frac{E_q}{2M(1 - \xi^2)} \right)^2 \leq \frac{q(x_1)q(x_2)}{1 - \xi^2}. \tag{13}$$

Taking into account that

$$t = \frac{4\xi^2 M^2 + |\Delta_{\perp}|^2}{1 - \xi^2} \tag{14}$$

one arrives at the result (1).

Note that excluding E_q from the bound (1) we obtain a constraint on H_q

$$|H_q(x, \xi, t)| \leq \sqrt{\left(1 - \frac{t_0}{t}\right)^{-1} \frac{q(x_1)q(x_2)}{1 - \xi^2}}. \tag{15}$$

Since $t \leq t_0 \leq 0$ this bound is weaker than the obsolete bound (6).

In a similar way one can derive the following bound for the gluon GPDs H_g , E_g (normalized according to the convention of ref. [12])

$$\begin{aligned}
& \left| H_g(x, \xi, t) - \frac{\xi^2}{1 - \xi^2} E_g(x, \xi, t) \right|^2 \\
& + \left| \frac{\sqrt{t_0 - t}}{2M\sqrt{1 - \xi^2}} E_g(x, \xi, t) \right|^2 \leq \frac{x^2 - \xi^2}{1 - \xi^2} g(x_1)g(x_2)
\end{aligned} \tag{16}$$

where $g(x)$ is the forward gluon distribution. In particular, this allows us to reproduce the following result from ref. [10]

$$\left| H_g(x, \xi, t) - \frac{\xi^2}{1 - \xi^2} E_g(x, \xi, t) \right| \leq \sqrt{\frac{x^2 - \xi^2}{1 - \xi^2} g(x_1)g(x_2)}. \tag{17}$$

By analogy with (15) we find from (16)

$$|H_g(x, \xi, t)| \leq \sqrt{\left(1 - \frac{t_0}{t}\right)^{-1} \frac{x^2 - \xi^2}{1 - \xi^2} g(x_1)g(x_2)}. \tag{18}$$

Although the inequalities derived here are stronger than the bounds on GPDs found earlier, one should not consider the new results as the best final estimates. Indeed, the derivation included the summation over quark polarization indices in the lhs of (7) and some information could be lost at this step.

I appreciate discussions with A. Belitsky, J.C. Collins, M. Diehl, N. Kivel, D. Müller, V.Yu. Petrov, M.V. Polyakov, A.V. Radyushkin, M. Strikman and O. Teryaev.

References

- [1] D. Müller, D. Robaschik, B. Geyer, F.-M. Dittes, and J. Hořejši, Fortschr. Phys. **42** (1994) 101.
- [2] A. V. Radyushkin, Phys. Lett. **B385** (1996) 333.
- [3] X. Ji, Phys. Rev. Lett. **78** (1997) 610; Phys. Rev. **D55** (1997) 7114.
- [4] J. C. Collins, L. Frankfurt, and M. Strikman, Phys. Rev. **D56** (1997) 2982.
- [5] A. V. Radyushkin, Phys. Rev. **D56** (1997) 5524.
- [6] A. D. Martin and M. G. Ryskin, Phys. Rev. **D57** (1998) 6692.
- [7] A. V. Radyushkin, Phys. Rev. **D59** (1999) 014030.
- [8] B. Pire, J. Soffer, O. Teryaev, Eur. Phys. J. **C8** (1999) 103.
- [9] X. Ji, J. Phys. **G24** (1998) 1181.
- [10] M. Diehl, T. Feldmann, R. Jakob, and P. Kroll, Nucl. Phys. **B596** (2001) 33.
- [11] M. Diehl (private communication).
- [12] M. Diehl, Eur. Phys. J. **C19** (2001) 485.